

Rapid Assessment Reference Condition Model

The Rapid Assessment is a component of the LANDFIRE project. Reference condition models for the Rapid Assessment were created through a series of expert workshops and a peer-review process in 2004-2005. For more information, please visit www.landfire.gov. Please direct questions to helpdesk@landfire.gov.

Potential Natural Vegetation Group (PNVG):

R0SPFI

Upper Subalpine Spruce-Fir - Central Rockies

General Information

Contributors (additional contributors may be listed under "Model Evolution and Comments")

Modelers

Kathy Roche kroche@fs.fed.us

Reviewers

Bill Baker bakerwl@uwyo.edu
Dennis Knight dhknight@uwyo.edu
Bill Romme romme@cnr.colostate.edu
u

Vegetation Type

Forested

Dominant Species*

PIEN
ABLA
PICO

General Model Sources

- Literature
 Local Data
 Expert Estimate

LANDFIRE Mapping Zones

10	21
19	22
20	29

Rapid Assessment Model Zones

- | | |
|--|--|
| <input type="checkbox"/> California | <input type="checkbox"/> Pacific Northwest |
| <input type="checkbox"/> Great Basin | <input type="checkbox"/> South Central |
| <input type="checkbox"/> Great Lakes | <input type="checkbox"/> Southeast |
| <input type="checkbox"/> Northeast | <input type="checkbox"/> S. Appalachians |
| <input type="checkbox"/> Northern Plains | <input type="checkbox"/> Southwest |
| <input checked="" type="checkbox"/> N-Cent.Rockies | |

Geographic Range

Central Rockies (Wyoming).

Biophysical Site Description

Upper subalpine zone in the central Rockies.

Vegetation Description

Engelmann spruce and subalpine fir dominate on most aspects with lodgepole pine comprising a greater component on dryer sites or earlier successional stages. *Vaccinium scoparium* is a common understory associate.

Disturbance Description

Fire Regime Group V is most likely, primarily long-interval (e.g.200+ year) stand replacement fires. In some areas, spruce beetle can influence successional stage, species composition and stand density. Spruce beetle may act to accelerate succession.

Adjacency or Identification Concerns

Adjacent to lower subalpine forests (Lodgepole-spruce-fir) and to krummholz, subalpine meadows, and alpine tundra types. Has more precipitation and longer winters than lower subalpine types.

Climate (severely dry conditions) is the primary driver of fire regimes in this system. Long-term changes in climate as well as interannual climate variability will affect the frequency of fire in this system.

This PNVG may be similar to the PNVGs R#SPFI from the Pacific Northwest model zone, R2SPFI from the Great Basin model zone, and R3SPFI from the Southwest model zone.

*Dominant and Indicator Species are from the NRCS PLANTS database. To check a species code, please visit <http://plants.usda.gov>.

Scale Description

Sources of Scale Data Literature Local Data Expert Estimate

Fires could range from 1,000's to 10,000's of acres. Variability of climate, topography and other site factors can result in a wide range of representation of successional stages on the landscape. Equilibrium landscapes are not likely to develop in areas <500,000 acres.

Issues/Problems

Model Evolution and Comments

Workshop code was USAL3.

Peer review was incorporated on 4/30/2005. Reviewers agreed that the model parameters were satisfactory and slight modifications were made to enhance the description.

Succession Classes

Succession classes are the equivalent of "Vegetation Fuel Classes" as defined in the Interagency FRCC Guidebook (www.frcc.gov).

Class A 15%

Early1 PostRep

Description

Early succession stage after long interval replacement fires. There can be extended periods (as long as 300 years) of grass/seedling stage after fire replacement events. This stage may occupy 3-50% of the landscape depending upon climatic conditions and variability of fire return intervals

Indicator Species* and Canopy Position

PIEN
PICO
ABLA

Upper Layer Lifeform

- Herbaceous
 Shrub
 Tree

Fuel Model no data

Structure Data (for upper layer lifeform)

	Min	Max
Cover	0 %	100 %
Height	no data	no data
Tree Size Class	no data	

- Upper layer lifeform differs from dominant lifeform. Height and cover of dominant lifeform are:

Class B 25%

Mid1 Closed

Description

High density saplings to poles. May occupy 5-50% of the landscape.

Indicator Species* and Canopy Position

PIEN
PICO
ABLA

Upper Layer Lifeform

- Herbaceous
 Shrub
 Tree

Fuel Model no data

Structure Data (for upper layer lifeform)

	Min	Max
Cover	40 %	100 %
Height	no data	no data
Tree Size Class	no data	

- Upper layer lifeform differs from dominant lifeform. Height and cover of dominant lifeform are:

Class C 20%

Late1 Open
Description

Low density saplings to poles. Primarily occurs after insects, disease or weather stress thins denser stands. This occupies 3-50% of landscape.

Indicator Species* and Canopy Position

PIEN
PICO
ABLA

Upper Layer Lifeform

- Herbaceous
- Shrub
- Tree

Fuel Model no data

Structure Data (for upper layer lifeform)

	Min	Max
Cover	0 %	40 %
Height	no data	no data
Tree Size Class	no data	

Upper layer lifeform differs from dominant lifeform. Height and cover of dominant lifeform are:

Class D 40%

Late1 Closed
Description

Pole- to larger diameter trees. This stage occupies 15 to 50% of the landscape.

Indicator Species* and Canopy Position

PIEN
ABLA

Upper Layer Lifeform

- Herbaceous
- Shrub
- Tree

Fuel Model no data

Structure Data (for upper layer lifeform)

	Min	Max
Cover	40 %	100 %
Height	no data	no data
Tree Size Class	no data	

Upper layer lifeform differs from dominant lifeform. Height and cover of dominant lifeform are:

Class E 0%

Late1 Closed
Description

Indicator Species* and Canopy Position

Structure Data (for upper layer lifeform)

	Min	Max
Cover	%	%
Height	no data	no data
Tree Size Class	no data	

Upper Layer Lifeform

- Herbaceous
- Shrub
- Tree

Fuel Model no data

Upper layer lifeform differs from dominant lifeform. Height and cover of dominant lifeform are:

Disturbances

Non-Fire Disturbances Modeled

- Insects/Disease
- Wind/Weather/Stress
- Native Grazing
- Competition
- Other:
- Other:

Fire Regime Group: 5

- I: 0-35 year frequency, low and mixed severity
- II: 0-35 year frequency, replacement severity
- III: 35-200 year frequency, low and mixed severity
- IV: 35-200 year frequency, replacement severity
- V: 200+ year frequency, replacement severity

*Dominant and Indicator Species are from the NRCS PLANTS database. To check a species code, please visit <http://plants.usda.gov>.

Historical Fire Size (acres)

Avg:
Min:
Max:

Fire Intervals (FI):

Fire interval is expressed in years for each fire severity class and for all types of fire combined (All Fires). Average FI is the central tendency modeled. Minimum and maximum show the relative range of fire intervals, if known. Probability is the inverse of fire interval in years and is used in reference condition modeling. Percent of all fires is the percent of all fires in that severity class. All values are estimates and not precise.

Sources of Fire Regime Data

- Literature
- Local Data
- Expert Estimate

	Avg FI	Min FI	Max FI	Probability	Percent of All Fires
Replacement	300	100	600	0.00333	99
Mixed					
Surface					
All Fires	300			0.00335	

References

Alexander R.R., G.R. Hoffman and J.M Wirsing. 1986. Forest Vegetation of the Medicine Bow National Forest in Southeastern Wyoming: A Habitat Type Classification. Research Paper RM -271, USDA FS RMRS Fort Collins, CO

Alexander, R.R. 1986. Silvicultural Systems and Cutting Methods for Old-Growth Spruce-Fir Forests in the Central and Southern Rocky Mountains. General Technical Report RM-126. Rocky Mountain Forest and Range Experiment Station, Fort Collins, CO

Alexander, R.R. 1988. Forest Vegetation on National Forests in the Rocky Mountain and Intermountain Regions: Habitat Types and Community Types. General Technical Report RM-162. USDA FS, RMRS, Fort Collins, CO.

Alexander, R.R. and O. Engelby. 1983. Engelmann spruce - subalpine fir. In: Silvicultural systems for the major forest types of the United States. Agriculture Handbook 445. Washington, D.C, U.S. Dept. of Agriculture.

Aplet, Gregory H.; Laven, Rod D.; Smith, F.W. 1988. Patterns of community dynamics in Colorado Engelmann spruce and subalpine fir forests. Ecology 69:312-319.

Arno, Stephen F. 2000. Fire in western forest ecosystems. In: Brown, James K.; Smith, Jane Kapler, eds. Wildland fire in ecosystems: Effects of fire on flora. Gen. Tech. Rep. RMRS-GTR-42-vol. 2. Ogden, UT: U.S. Department of Agriculture, Forest Service, Rocky Mountain Research Station: 97-120

Baker, W. L. and R. Knight. 2000. Roads and forest fragmentation in the Southern Rocky Mountains. In Forest Fragmentation in the Southern Rocky Mountains. R. Knight, F.W. Smith, W.H. Romme and W.L. Baker eds. University Press of Colorado, Boulder, Colorado: 97-122.

Baker, W.L. 2000. Measuring and Analyzing Forest Fragmentation in the Rocky Mountains and Western United States. In Forest Fragmentation in the Southern Rocky Mountains. R. Knight, F.W. Smith, W.H. Romme and W.L. Baker eds. University Press of Colorado, Boulder, Colorado: 55-94.

Baker, W.L. and K. F. Kipfmueller. 2001. Spatial Ecology of Pre-Euro-American Fires in a Southern Rocky Mountain Subalpine Forest Landscape. The Professional Geographer. Vol 53 Number 2 pp 248-262.

Baker, W.L. and T.T. Veblen. 1990. Spruce Beetles and Fires in the Nineteenth-Century Subalpine forests of

*Dominant and Indicator Species are from the NRCS PLANTS database. To check a species code, please visit <http://plants.usda.gov>.

Western Colorado, U.S.A. *Arctic and Alpine Research*, Vol 22, No 1, 1990, pages 65-80.

Baker, William L. 1994. *Landscape Structure Measurements for Watersheds in the Medicine Bow National Forest Using GIS Analysis* Department of Geography and Recreation, Univ. of Wyoming prepared under agreement with the USDA FS MBNF. On file at Medicine Bow-Routt NFs and Thunder Basin NG Supervisor's Office, Laramie, WY.

Barrett, Stephen W. 1994. Fire regimes on andesitic mountain terrain in northeastern Yellowstone National Park. *International Journal of Wildland Fire* 4: 65-76.

Buechling, A., and W. L. Baker. 2004. A fire history from tree rings in a high-elevation forest of Rocky Mountain National Park. *Canadian Journal of Forest Research* 34: 1259-1273.

Buskirk, S.W., W.H. Romme, F.W. Smith, and R. Knight. 2000. An Overview of Forest Fragmentation in the Southern Rocky Mountains. In *Forest Fragmentation in the Southern Rocky Mountains*. R. Knight, F.W. Smith, W.H. Romme and W.L. Baker eds. University Press of Colorado, Boulder, Colorado: 3-14.

Clagg, H.B. 1975. *Fire ecology in high-elevation forests in Colorado*. M.S. Thesis, Colorado State University, Fort Collins, Colorado.

Coleman, M.D., T.M. Hinckley, G. McNaughton and B.A. Smit. 1992. Root cold Hardiness and Native Distribution of Sub-alpine Conifers, *Canadian Journal of Forest Research* Vol. 22 no 7, (July 1992) p. 932-938.

Crane, M.F. 1982. *Fire Ecology of Rocky Mountain Region Forest Habitat Types*. Report prepared under contract to USDA FS Region 2.

Despain, D.G. 1973. *Vegetation of the Big Horn Mountains in Relation to Substrate and Climate*. *Ecological Monographs* 43:329-355.

Despain, D.G. and R.E. Sellers. 1977. *Natural Fire in Yellowstone National Park*. *Western Wildlands*, summer 1977.

Dillon, G. K., D. Knight and C. Meyer. 2003. *Historic Variability for Upland Vegetation in the Medicine Bow National Forest*. Department of Botany, Univ. of Wyoming: prepared under agreement with the USDA FS MBNF 1102-0003-98-043

Graham, R.T. A.E. Harvey, M.F. Jurgensen, T.B. Jain, J.R. Tonn and D.S. Page-Dumrose. 1994. *Managing Coarse Woody Debris in Forests of the Rocky Mountains*. Research Paper INT-RP-477. USDA FS, Intermountain Research Station (now RMRS), Fort Collins, CO.

Griggs, R.F. 1938. Timberlines in the Northern Rocky Mountains. *Ecology* volume 19, number 4 (October 1938) pages 548-564.

Griggs, R.F. 1946. The timberlines of Northern America and their Interpretation. *Ecology* volume 27 number 4 (October 1946) pages 275-289.

Hinds, T.E., F.G. Hawksworth and R.W. Davidson. 1965. Beetle-killed Engelmann Spruce: Its deterioration in Colorado. *J. For.* 63(7): 536-542.

Jenkins, M. J., C. A. Dicus and E. G. Hebertson. 1998. Post-fire succession and disturbance interactions on an intermountain subalpine spruce-fir forest. Pp. 219-229 in Pruden, T. L. and L. A. Brennan, eds. Proceedings, Symposium: Fire in Ecosystem Management: Shifting the paradigm from suppression to prescription. Tall Timbers Fire Ecology Conference Proceedings, No. 20. Tall Timbers Research Station, Tallahassee, FL.

Jones, G. P., and S. M. Ogle. 2000. Characterization abstracts for vegetation types on the Bighorn, Medicine Bow, and Shoshone National Forests. Laramie. Prepared for USDA, Forest Service, Region 2, by George Jones and Steve Ogle, WYNDD, UW, Laramie WY.

Kane, T.L., B.G. Brown, R. Sharman. 1999. A Preliminary Climatology of Upper Level Turbulence Reports. Preprints, 8th Conf. on Aviation, Range and Aerospace Meteorology, 10-15 January, Dallas, TX, American Meteorology Society, 363-367.

Kipfmueller, K. F. and W. L. Baker. 2000. A fire-history of a subalpine forest in south-eastern Wyoming, USA. *Journal of Biogeography* 27 pages 71-85.

Kipfmueller, K.F. 1997. A fire history of a subalpine forest in southeastern Wyoming. Thesis. University of Wyoming. Laramie, WY.

Kipfmueller, K.F. and W.L. Baker. 1998a. A comparison of three techniques to date stand-replacing fires in lodgepole pine forests. *Forest Ecology and Management* 104(1998) 171-177.

Kipfmueller, Kurt F.; Baker, William L. 2000. A fire history of a subalpine forest in southeastern Wyoming, USA. *Journal of Biogeography* 27:71-85.

Knight, D.H. 1987. Ecosystem Studies in the Subalpine Coniferous Forests of Wyoming. In: Management of Subalpine Forests: Building on 50 years of Research: Proceedings of a Technical Conference. General Technical Report RM-149. USDA Forest Service Rocky Mountain Forest and Range Experiment Station. Fort Collins, CO.

Knight, D.H. 1994. Mountains and Plains, The Ecology of Wyoming Landscapes. Yale University Press, New Haven, CT.

Knight, D.H. and W.A. Reiners. 2000. Natural patterns in southern Rocky Mountain landscapes and their relevance to forest management. In *Forest Fragmentation in the Southern Rocky Mountains*. R. Knight, F.W. Smith, W.H. Romme and W.L. Baker eds. University Press of Colorado, Boulder, Colorado:15-30.

Knight, D.H., A.D. Anderson, G.T. Baxter, K.L. Diem, M. Parker, P.A. Rechar, P.C. Singleton, J.F. Thilenius, A.L. Ward and R.W. Weeks. 1975. Final Report the Medicine Bow Ecology Project, The potential Sensitivity of Various Ecosystem Components to Winter Precipitation Management in The Medicine Bow Mountains, Wyoming. Prepared for the Division of Atmospheric Water Resources Management, Bureau of Reclamation, USDI, Denver, CO by the Rocky Mountain Forest and Range Experiment Station, USFS and the Wyoming Water Resource Research Institute.

Logan, J.A., J.M. Schmid and M.S. Mehl. 1980. A Computer Program to Calculate Susceptibility of Spruce-Fir Stands to Spruce Beetle Outbreaks. USDA Forest Service Research Note RM-303. Rocky Mountain Forest and Range Experiment Station. Fort Collins, CO.

- Loope, Lloyd L.; Gruell, George E. 1973. The ecological role of fire in the Jackson Hole area, northwestern Wyoming. *Quaternary Research* 3(3): 425-443.
- Mehl, M. 1992. Old-Growth Descriptions for the Major Forest Cover Types in the Rocky Mountain Region. In: *Old-Growth Forests in the Southwest and Rocky Mountain Regions Proceedings of A Workshop*, Rocky Mountain Forest and Range Exp. Sta. Gen. Tech. Report RM-213, USDA FS RMRMS Fort Collins, CO.
- Merrill, E.H., T.W. Kohley, M.E. Herdendorf, W.A. Reiners, K.L. Driese, R.W. Marrs, S.A. Anderson. 1996. Wyoming GAP Analysis Project Final Report. University of Wyoming Department of Physiology and Department of Botany, Wyoming Cooperative Fish and Wildlife Research Unit and USGS Biological Resources Division. From <http://www.sdvc.uwyo.edu/wbn/abstract.html>
- Meyer, C. B., and D. H. Knight. 2001. Historic variability of upland vegetation in the Bighorn National Forest, Wyoming. Draft report, November 30, 2001.
- Mielke, J.L. 1950. Rate of Deterioration of beetle-killed Engelmann spruce. *J. For.* 48(12): 882-888.
- Moir, W.H. Ecological Concepts in Old-Growth Forest Definition. In: *Old-Growth Forests in the Southwest and Rocky Mountain Regions Proceedings of A Workshop*, Rocky Mountain Forest and Range Exp. Sta. Gen. Tech. Report RM-213, USDA FS RMRMS Fort Collins, CO.
- Pennanen, J. 2002. Forest Age Distribution Under Mixed-Severity Fire Regimes – A simulation-based analysis for middle boreal Fennoscandia. *Silva Fennica: quarterly issues*: 36(1): 213-231.
- Schmid, J. M., and R. H. Frye. 1977. Spruce beetle in the Rockies. *Spruce Beetle in the Rockies*. General Technical Report RM 49. USDA FS Rocky Mountain Forest and Range Experiment Station, Fort Collins, CO. 38 pp.
- Schmid, J.M. and R.C. Beckwith. 1977. The Spruce Beetle. Pest Leaflet 127. USDA Forest Service 7 pages.
- Schmid, J.M. and R.H. Frye. 1976. Stand Ratings for Spruce Beetles. USDA Forest Service Research Note RM-309. Rocky Mountain Forest and Range Experiment Station. Fort Collins, CO.
- Schmid, J.M. and S.A. Mata. 1996. Natural Variability of Specific Forest Insect Populations and Their Associated Effects in Colorado. USDA FS Rocky Mountain Forest and Range Experiment Station General Technical Report RM-GTR-275, RMF&R Experiment Station, Fort, Collins, CO.
- Schmid, J.M. and T.E. Hinds. 1974. Development of Spruce-Fir Stands Following Spruce Beetle Outbreaks. Research Paper RM-131, USDA Forest Service, Rocky Mountain Forest and Range Experiment Station. Fort Collins, CO.
- Schmidt, Kirsten M, Menakis, James P., Hardy, Colin C., Hann, Wendel J., Bunnell, David L. 2002. Development of coarse-scale spatial data for wildland fire and fuel management. Gen. Tech. Rep. RMRS-GTR-87. Fort Collins, CO: U.S. Department of Agriculture, Forest Service, Rocky Mountain Research Station. 41 p. + CD.
- Schrupp, D.L., W.A. Reiners, T.G. Thompson, L.E. O'Brien, J.A. Kindler, M.B. Wunder, J.F. Lowsky, J.C. Buoy, L. Satcowitz, A.L. Cade, J.D. Stark, K.L. Driese, T.W. Owens, S.J. Russo, and F. D'Erchia. 2000.

Colorado Gap Analysis Program: A Geographical Approach to Planning for Biological Diversity - Final Report, USGS Biological Resource Division, Gap Analysis Program and Colorado Division of Wildlife, Denver, CO.

Sherriff, R.; Veblen, T.T.; Sibold, J.S. 2001. Fire history in high elevation subalpine forests in the Colorado Front Range. *Ecoscience* 8:369-380.

Sibold, J. 2001. The forest fire regime of an upper montane and subalpine forest, Wild Basin, Rocky Mountain National Park. M.S. Thesis, University of Colorado, Boulder, CO.

Stahelin, R. 1943. Factors Influencing the Natural Restocking of High Altitude Burns by Coniferous Trees in the Central Rocky Mountains. *Ecology* 24:19-30.

Veblen, T. T., K. S. Hadley and M. S. Reid. 1991. Disturbance and stand development of a Colorado subalpine forest. *Journal of Biogeography* (1991) 18:707-716.

Veblen, T. T., K. S. Hadley, E. M. Nel, T. Kitzberger, M. S. Reid, and R. Villalba. 1994. Disturbance regime and disturbance interactions in a Rocky Mountain subalpine forest. *Journal of Ecology* 82:125-135.

Veblen, T. T., K. S. Hadley, M. S. Reid and A.J. Rebertus. 1989. Blowdown and stand development in a Colorado subalpine forest. *Canadian Journal of Forest Resources*. Vol 19: 1218-1225.

Veblen, T.T.; Kitzberger, T. 2002. Inter-hemispheric comparison of fire history: The Colorado Front Range, U.S.A. and the Northern Patagonian Andes, Argentina. *Plant Ecology*, in press.

Whipple, S.A.; Dix, R.L. 1979. Age structure and successional dynamics of a Colorado subalpine forest. *American Midland Naturalist* 101:142-158.